

Heart Vessels (2012) 27:258–264
DOI 10.1007/s00380-011-0146-0

ORIGINAL ARTICLE

Off-pump surgery for the poor ventricle?

Maximilian Y. Emmert · Sacha P. Salzberg · Oliver M. Theusinger · Hector Rodriguez · Simon H. Sündermann · Andre Plass · Christoph T. Starck · Burkhardt Seifert · Werner Baulig · Simon P. Hoerstrup · Stephan Jacobs · Jürg Grünenfelder · Volkmär Falk

Received: 6 January 2011 / Accepted: 1 April 2011 / Published online: 17 May 2011
© Springer 2011

Abstract Severely decreased ejection-fraction is an established risk-factor for worse outcome after cardiac surgery. We compare outcomes of off-pump coronary artery bypass grafting (OPCAB) and on-pump CABG (ONCABG) in patients with severely compromised EF. From 2004 to 2009, 478 patients with a decreased EF $\leq 35\%$ underwent myocardial-revascularization. Patients received either OPCAB ($n = 256$) or ONCABG ($n = 222$). Propensity score (PS), including 50 preoperative risk-factors, was used to balance characteristics between groups. PS adjusted logistic regression analysis was performed to assess mortality and major adverse cardiac and cerebrovascular events (MACCE). A composite endpoint for major non-cardiac complications such as respiratory failure, renal failure, rethoracotomy was applied. Complete revascularization (CR) was assumed when the number of distal anastomoses was larger than that of diseased vessels. There was no difference for mortality (2.3 vs. 4.1%; PS-adjusted odds ratio (PS-OR) = 1.05; $p = 0.93$) and MACCE (13.7 vs. 17.6%; PS-OR = 1.22; $p = 0.50$) including myocardial-infarction (1.4 vs. 4.9%; PS-OR = 0.39; $p = 0.26$), low cardiac output (2.3 vs. 4.7%; PS-OR = 0.75; $p = 0.72$) and stroke (2.3 vs.

2.7%; PS-OR = 0.69; $p = 0.66$). OPCAB patients presented with a trend to less frequent occurrence of the non-cardiac composite (12.1 vs. 22.1%; PS-OR = 0.54; $p = 0.059$) including renal dysfunction (PAOR = 0.77; 95% CI 0.31–1.9; $p = 0.57$), bleeding (PAOR = 0.42; 95% CI 0.14–1.20; $p = 0.10$) and respiratory failure (PAOR = 0.39; 95% CI 0.05–3.29; $p = 0.39$). The rate of complete revascularization was similar (92.2 vs. 92.8%; PS-OR = 0.75; $p = 0.50$). OPCAB in patients with severely decreased EF is safe and feasible. It may even benefit these patients in regard to non-cardiac complications and does not come at cost of less complete revascularization.

Keywords Ejection fraction · Off-pump surgery · Coronary artery bypass grafting · Heart failure

Introduction

Surgical revascularization for patients with severe left ventricular dysfunction remains a challenge. Various studies report an increased peri-operative mortality and morbidity for these patients [1, 2]. In the EuroScore risk stratification system (<http://www.euroscore.org>), decreased ventricular function is an independent risk factor for worse outcome after cardiac surgery.

Conventional CABG is performed with cardio-pulmonary bypass (CPB), which is associated with complications such as renal dysfunction, stroke, and other neurological complications [3] as well as an increased risk for a systemic inflammatory response syndrome (SIRS) [4]. The compromised outcomes for patients undergoing CABG with ventricular dysfunction may be in part linked to the use of CPB and the application of cardioplegic solutions to achieve cardiac arrest [4, 5].

M. Y. Emmert (✉) · S. P. Salzberg · H. Rodriguez · S. H. Sündermann · A. Plass · C. T. Starck · S. P. Hoerstrup · S. Jacobs · J. Grünenfelder · V. Falk
Department of Cardiovascular Surgery, University Hospital Zurich, Raemi Street 100, 8091 Zurich, Switzerland
e-mail: maximilian.emmert@web.de

O. M. Theusinger · W. Baulig
Institute for Anaesthesiology, University Hospital Zurich, Zurich, Switzerland

B. Seifert
Biostatistics Unit, Institute of Social and Preventive Medicine, University of Zurich, Zurich, Switzerland

Off-pump coronary artery bypass surgery (OPCAB) has a comparable risk-adjusted mortality and morbidity as on-pump CABG [6–8]. Current reports suggest OPCAB to be superior for high-risk patients [9]. For patients with severe ventricular dysfunction, only few reports are available [10–13].

In this study, we evaluate safety and feasibility of OPCAB surgery in patients with a decreased ejection fraction ($\leq 35\%$).

Materials and methods

From 2004 to 2009, 3,981 patients underwent surgical revascularization at our institution. Of these patients, 478 presented with severe ventricular dysfunction and an ejection fraction $\leq 35\%$. Of these, 53% ($n = 256$) received off-pump surgery (OPCAB) and 47% ($n = 222$) conventional CABG. All data were collected prospectively and the study was approved by an institutional review board (IRB), including a waiver of informed consent. Indication for surgery was: elective (66.8 vs. 64.4%; $p = 0.63$), urgent (32.4 vs. 27.7%; $p = 0.27$), and emergent (5.5 vs. 3.2%; $p = 0.26$) respectively. The EuroScore (<http://www.EuroScore.org>) was used for preoperative risk stratification. The ejection fraction was documented by preoperative transthoracic echocardiography and was confirmed with in a left ventriculogram, which was obtained during coronary angiography to define the target vessels. In addition, these data were correlated to intraoperative trans-esophageal echocardiography (TOE) in all patients as TOE is part of our OPCAB practice (Table 1).

Surgical technique

CABG was performed with administration of standardized cardio-pulmonary bypass techniques and proximal anastomosis was done with cross-clamping of the aorta. OPCAB procedures were performed as previously described [14]. Briefly, heparin was given to achieve an active clotting time (ACT) in excess of 350 s. Pacemaker wires were placed epicardially, before a stabilizer (Octopus® 4 Tissue Stabilizer, Medtronic, Minneapolis, USA) was applied for target vessel exposure. Shunt insertion was (ClearView® Intracoronary Shunt, Medtronic, Minneapolis, USA) was routinely performed to maintain distal perfusion during anastomosis and a mister blower (Guidant, Indianapolis, USA) with CO₂ and water was used to clear the surgical field. If no T-Graft was performed and whenever a proximal anastomosis was necessary, it was performed in ‘no touch’ fashion using the heartstring device (HEART-STRING™ Proximal Seal System, Guidant, Indianapolis, USA). After LIMA to LAD grafting, the right coronary

Table 1 Preoperative characteristics and demographics

| Parameter | OPCAB <i>n</i> = 256 | CABG <i>n</i> = 222 | <i>p</i> value |
|--------------------------|-------------------------|------------------------|----------------|
| Age (years) | 64 ± 10 | 63 ± 9 | 0.27 |
| Male (%) | 81 | 85 | 0.22 |
| Female (%) | 19 | 15 | 0.23 |
| EuroScore | 4.6 ± 1.0 | 4.9 ± 1.0 | 0.06 |
| BMI (kg/m ²) | 27 ± 4 | 27 ± 4 | 0.54 |
| Elective (%) | 66.8 | 64.4 | 0.63 |
| Urgent (%) | 32.4 | 27.7 | 0.27 |
| Emergent (%) | 5.5 | 3.2 | 0.26 |
| Sinus rhythm (%) | 91.6 | 98.6 | 0.002 |
| Atrial fibrillation (%) | 4.9 | 0.9 | 0.03 |
| Pacemaker (%) | 3.5 | 0.5 | 0.04 |
| No. of diseased vessels | 2.79 ± 0.45 | 2.95 ± 0.23 | 0.001 |
| 1-vessel disease (%) | 1.5 | 2.3 | 0.74 |
| 2-vessel disease (%) | 18.4 | 11.7 | 0.56 |
| 3-vessel disease (%) | 80.1 | 86.0 | 0.09 |
| Left main disease (%) | 28.5 | 23.9 | 0.26 |
| CCS 1 (%) | 7.6 | 7.2 | 1.00 |
| CCS 2 (%) | 29.4 | 28.4 | 0.83 |
| CCS 3 (%) | 42.2 | 36.9 | 0.28 |
| CCS 4 (%) | 20.9 | 27.5 | 0.11 |
| NYHA 1 (%) | 51.5 | 49.5 | 0.70 |
| NYHA 2 (%) | 26.4 | 22.1 | 0.32 |
| NYHA 3 (%) | 19.5 | 17.1 | 0.55 |
| NYHA 4 (%) | 2.6 | 11.3 | 0.001 |
| Redo surgery (%) | 5.5 | 4.5 | 0.68 |

EF ejection fraction, *BMI* body mass index, *CCS* Canadian Cardiovascular Society Angina Classification, *NYHA* New York Heart Association

system was approached, and finally the circumflex territory was addressed. Standardized ultrasound flow measurement (MediStim QuickFit®) was carried out in all patients [15].

Completeness of revascularization

The anticipated complete revascularization was based on the defined target vessels reported on the preoperative angiography. This information was compared to the performed grafts documented in the postoperative surgical report and was assumed to be complete when the total number of distal grafts was higher than that of diseased vessels reported on the preoperative angiography [7].

Statistical analysis

Endpoints analyzed are mortality and major adverse cardiac and cerebrovascular events (MACCE). A composite endpoint including major non-cardiac adverse events such

as respiratory failure, renal failure, and rethoracotomy for bleeding was created. Crossover patients were analyzed on an ‘*Intention-to-treat*’ basis.

Continuous data are presented as mean \pm standard deviation and are compared using the Mann–Whitney test. Categorical data are presented as number and percentage and are compared using the Chi-square test or Fisher’s exact test where appropriate. Odds ratios with 95% confidence intervals are computed using univariate logistic regression. A propensity score (PS) was calculated using logistic regression with preoperative variables to balance characteristics between both groups. In case of missing values in preoperative variables were replaced using regression methods. The PS then was divided into quintiles and analyzed as a categorical variable. PS-adjusted logistic regression analysis was performed to assess binary endpoints and two-way analysis of variance for continuous endpoints.

The data-set of preoperative variables for the PS score included preoperative patient characteristics such as cardiovascular risk-factors and co-morbidities including cerebral-vascular events, peripheral artery disease (PAD), chronic-obstructive pulmonary-disease (COPD), and renal disease. These parameters were defined in accordance with the definitions of the EuroScore Risk Stratification system (<http://www.euroscore.org>); Cerebrovascular disease defined as the history of transient ischemic attacks (TIA) or stroke, COPD defined with the long-term use of bronchodilators or steroids for lung disease, renal disease as creatinine levels $>200 \mu\text{mol/l}$ and PAD defined as one or more of claudication, carotid occlusion or $>50\%$

stenosis, previous or planned intervention on the abdominal aorta, limb arteries or carotids. Diabetes mellitus was defined as type 2 insulin-dependant diabetes (Table 2).

Cardiac-related preoperative conditions were: preceding myocardial-infarction (MI), recent MI within 3 months prior to surgery, preceding cardiogenic-shock, congestive heart-failure, instable angina pectoris, arrhythmias, number of diseased coronary-vessels, previous CABG, elective, urgent/or emergent presentation, previous PTCA, previous stent implantation, NYHA class, CCS class, logistic EuroScore and others. All analyses were performed using SPSS 18 (SPSS Inc., Chicago, IL, USA). p values <0.05 are assumed to be statistically significant.

Results

Patient demographics

OPCAB patients and CABG patients were comparable with regards to mean age, gender, and EuroScore. In brief, OPCAB patients presented with significantly more peripheral artery disease (18.3 vs. 11.3%; $p = 0.039$) and diabetes (35.2 vs. 22.5%; $p = 0.003$), whereas patients undergoing on-pump CABG suffered more frequently from chronic obstructive pulmonary disease (COPD) (4.7 vs. 10.4%; $p = 0.022$) and presented more frequently with instable angina (14.7 vs. 27.5%; $p = 0.005$). The number of patients presenting with significant left main disease (28.5 vs. 23.9%; $p = 0.26$) or for redo surgery (5.5 vs. 4.5%; $p = 0.68$) was comparable in both groups.

Table 2 Risk factors and co-morbidities

| Parameter | OPCAB $n = 256$ | CABG $n = 222$ | p value |
|--|-----------------|----------------|-----------|
| Hypercholesterolemia (%) | 65.0 | 71.6 | 0.20 |
| Hypertension (%) | 58.1 | 50.4 | 0.10 |
| Positive family history (%) | 32.0 | 32.0 | 1.00 |
| Diabetes | 35.2 | 22.5 | 0.003 |
| Smoking (%) | 61.0 | 59.5 | 0.76 |
| Adiposities (%) | 48.8 | 41.0 | 0.09 |
| PAD (%) | 18.3 | 11.3 | 0.039 |
| COPD (%) | 4.7 | 10.4 | 0.022 |
| Recent myocardial infarction (<90 days) (%) | 20.7 | 31.5 | 0.009 |
| Previous MI (>90 days) (%) | 48.0 | 59.9 | 0.01 |
| Previous cardiogenic shock (>90 days) (%) | 14.7 | 17.1 | 0.56 |
| Cerebrovascular disease (%) | 2.0 | 0.0 | 0.06 |
| Previous syncope (%) | 5.6 | 1.4 | 0.03 |
| Renal disease (%) | 4.2 | 4.5 | 1.00 |
| Instable angina (%) | 14.7 | 27.5 | 0.005 |
| IABP preoperative (%) | 18.8 | 25.2 | 0.09 |

PAD peripheral artery disease,
COPD chronic obstructive
pulmonary disease,
MI myocardial infarction,
IABP intra aortic balloon pump

Outcome data before and after propensity adjustment (Table 3)

OPCAB patients had a comparable mortality rate (2.3 vs. 4.1%; odds ratio (OR) = 0.57; 95% CI 0.20–1.62; $p = 0.29$) and similar MACCE (13.7 vs. 17.6%; OR = 0.74; 95% CI 0.44–1.22; $p = 0.24$) such as myocardial infarction (1.4 vs. 4.1%; OR = 0.34; 95% CI 0.71–1.58; $p = 0.17$) and low cardiac output (2.3 vs. 4.7%; OR = 0.48; 95% CI 0.11–2.23; $p = 0.35$). The composite of non-cardiac adverse events was lower in the OPCAB group (12.1 vs. 22.1%; OR = 0.49; 95% CI 0.30–0.80; $p = 0.004$). This was mainly driven by the reduced occurrence of bleeding complications (6.3 vs. 9.0%; OR = 0.67; 95% CI 0.34–1.33; $p = 0.26$), renal dysfunction (4.3% vs. 8.1%; OR = 0.50; 95% CI 0.23–1.10; $p = 0.08$) and respiratory failure (1.1 vs. 4.2%; OR = 0.27; 95% CI 0.33–2.12; $p = 0.21$). The beneficial effect of OPCAB in regard to respiratory complications was also reflected in a lower rate of pleural effusions and pneumothorax (3.5 vs. 7.5%; OR = 0.25; 95% CI 0.16–1.25; $p = 0.13$) as well as in the overall shorter time to extubation (<12 h) (58.9 vs. 36.9%; OR = 2.44; 95% CI 1.63–3.65; $p < 0.001$) indicating a more straight-forward postoperative course. Neurological outcomes including the occurrence of stroke were

comparable in both groups (2.3 vs. 2.7%; OR = 0.86; 95% CI 0.28–2.72; $p = 0.80$).

After propensity score adjustment, OPCAB patients still displayed comparable mortality (propensity-adjusted odds ratio (PAOR) = 1.05; 95% CI 0.30–3.63; $p = 0.93$) and MACCE (PAOR = 1.22; 95% CI 0.68–2.22; $p = 0.50$). In regard to the non-cardiac complications including renal dysfunction (PAOR = 0.77; 95% CI 0.31–1.9; $p = 0.57$), bleeding (PAOR = 0.42; 95% CI 0.14–1.20; $p = 0.10$) and respiratory failure (PAOR = 0.39; 95% CI 0.05–3.29; $p = 0.39$), the beneficial effect of OPCAB remained visible. Similarly, a trend towards less occurrence of the non-cardiac composite was still detectable for OPCAB, however, failed to achieve statistical significance after adjustment (PAOR = 0.54; 95% CI 0.29–1.02; $p = 0.059$).

Intra-operative data and completeness of revascularization (Table 4)

The need for intra-operative implantation of an IABP was similar in both groups (5.9 vs. 7.2%; $p = 0.32$). Conversion to CPB became necessary in 5.9% of all OPCAB patients. If converted, the operation was continued in beating-heart fashion.

Table 3 Crude outcome and propensity adjusted outcome

| Parameter | OPCAB <i>n</i> = 256 | CABG <i>n</i> = 222 | OR | CI 95% | <i>p</i> value | PA OR | PA CI 95% | PA <i>p</i> value |
|--------------------------------------|-------------------------|------------------------|------|------------|------------------|-------|------------|----------------------|
| Mortality (%) | 2.3 | 4.1 | 0.57 | 0.20–1.62 | 0.29 | 1.05 | 0.30–3.63 | 0.93 |
| Neurological events (central) (%) | 2.3 | 2.7 | 0.86 | 0.28–2.72 | 0.80 | 0.69 | 0.14–3.54 | 0.66 |
| Neurological events (peripheral) (%) | 1.2 | 0.5 | 2.62 | 0.27–25.37 | 0.40 | 1.14 | 0.07–18.84 | 0.93 |
| Re-thoracotomy for bleeding (%) | 6.3 | 9.0 | 0.67 | 0.34–1.33 | 0.26 | 0.42 | 0.14–1.20 | 0.10 |
| Myocardial infarction (%) | 1.4 | 4.1 | 0.34 | 0.71–1.58 | 0.17 | 0.39 | 0.07–2.01 | 0.26 |
| Low cardiac output (%) | 2.3 | 4.7 | 0.48 | 0.11–2.23 | 0.35 | 0.75 | 0.15–3.70 | 0.72 |
| Graft occlusion (%) | 0.0 | 1.9 | – | – | 0.32 | – | – | 0.32 |
| Cardiac tamponade (%) | 0.0 | 0.5 | – | – | 1.00 | – | – | 1.00 |
| Arrhythmia (%) | 6.9 | 3.7 | 1.91 | 0.64–5.67 | 0.24 | 2.15 | 0.63–7.30 | 0.22 |
| IABP postop (%) | 0.0 | 2.7 | – | – | 0.01 | – | – | 0.01 |
| Renal dysfunction (%) | 4.3 | 8.1 | 0.50 | 0.23–1.10 | 0.08 | 0.77 | 0.31–1.90 | 0.57 |
| No ventilation (%) | 1.0 | 0.9 | 1.12 | 0.16–7.99 | 0.91 | 6.96 | 0.91–53.11 | 0.61 |
| Ventilation <12 h (%) | 58.9 | 36.9 | 2.44 | 1.63–3.65 | <0.001 | 1.39 | 0.79–2.43 | 0.25 |
| Prolonged ventilation >24 h (%) | 8.3 | 11.7 | 0.69 | 0.36–1.33 | 0.27 | 1.59 | 0.70–3.42 | 0.27 |
| Respiratory failure (%) | 1.1 | 4.2 | 0.27 | 0.33–2.12 | 0.21 | 0.39 | 0.05–3.29 | 0.39 |
| Pleural effusions/pneumothorax (%) | 3.5 | 7.5 | 0.25 | 0.16–1.25 | 0.13 | 0.56 | 0.19–1.66 | 0.30 |
| Sinus rhythm (%) | 87.4 | 93.7 | 0.47 | 0.22–0.97 | 0.04 | 0.52 | 0.23–1.18 | 0.12 |
| Atrial fibrillation (%) | 9.1 | 5.9 | 1.60 | 0.72–3.57 | 0.24 | 1.90 | 0.80–4.48 | 0.15 |
| MACCE (%) | 13.7 | 17.6 | 0.74 | 0.45–1.22 | 0.24 | 1.22 | 0.68–2.20 | 0.50 |
| Non-cardiac composite (%) | 12.1 | 22.1 | 0.49 | 0.30–0.80 | 0.004 | 0.54 | 0.29–1.02 | 0.059 |

OR odds ratio, 95% CI confidence interval 95%, PA OR propensity adjusted OR, PA 95% CI propensity adjusted 95% CI, PA *p* value propensity adjusted *p* value, IABP Intra-aortic balloon pump, MACCE major adverse cardiac and cerebrovascular events

Table 4 Intraoperative data

| Parameter | OPCAB <i>n</i> = 256 | CABG <i>n</i> = 222 | <i>p</i> value |
|---------------------------------------|----------------------|---------------------|------------------|
| CPB conversion (%) | 5.9 | – | – |
| CPB time (min) | – | 107 ± 46 | – |
| Aortic x-clamp time (min) | – | 48 ± 27 | – |
| Arterial grafts per patient | 1.49 ± 0.98 | 1.35 ± 0.80 | 0.28 |
| LIMA (%) | 96.9 | 95.0 | 0.67 |
| RIMA (%) | 30.9 | 26.6 | 0.73 |
| Radial artery (%) | 16.0 | 3.6 | <0.001 |
| SVG per patient | 1.93 ± 1.28 | 2.50 ± 1.10 | <0.001 |
| Use of SVG (%) | 81.6 | 89.2 | 0.037 |
| Total number of proximal anastomoses | 1.12 ± 0.61 | 1.44 ± 0.64 | 0.05 |
| Total number of grafts per patient | 3.42 ± 1.05 | 3.85 ± 0.81 | <0.001 |
| No of diseased vessels | 2.79 ± 0.45 | 2.95 ± 0.23 | 0.001 |
| Completeness of revascularization (%) | 92.2 | 92.8 | 0.50 |
| IABP intraoperative (%) | 5.9 | 7.2 | 0.32 |

OPCAB patients presented with a lower mean number of the diseased coronary vessels (2.79 ± 0.45 vs. 2.95 ± 0.23 ; $p = 0.001$) and also received a lower number of total distal grafts (3.42 ± 1.05 vs. 3.85 ± 0.81 ; $p < 0.001$). For similar proportions in both groups, revascularization was possible without necessity of any proximal anastomosis (9.8 vs. 10.8%; $p = 0.42$). Complete revascularization was achieved in similar levels for both groups (92.2 vs. 92.8%; $p = 0.50$).

Discussion

OPCAB is safe and feasible in high-risk patients with severely decreased EF requiring myocardial revascularization. Mortality and MACCE are comparable to patients undergoing on-pump surgery. Furthermore, our results highlight that OPCAB patients presented with a trend to less non-cardiac complications, including renal dysfunction and postoperative bleeding. These patients also appeared to have decreased respiratory complications, which was also reflected in an overall faster extubation indicating a straighter postoperative course.

Although a recent prospective randomized trial identified OPCAB to have no advantage in low-risk patients [16], it appears to be the superior method of revascularization in high-risk patients [9, 17, 18].

Our results are in line with Stamou et al. [17] who recently reviewed 513 high-risk patients. Of these patients, 228 presented with an EF $<34\%$ either undergoing on-pump CABG ($n = 102$) or off-pump CABG ($n = 126$). They identified OPCAB to have a lower mortality and a comparable event-free survival. The authors concluded that off-pump CABG can be performed with a reasonably low morbidity and lower mortality in high-risk patients. In

comparison to our study, Stamou et al. [17] did not specifically focus on the outcome of patients with low EF, but on patients with a generally high-risk profile, including factors such as renal-failure, recent myocardial-infarction, cerebrovascular disease and advanced age.

Among 1,398 evaluated high-risk patients, Al-Ruzzeh et al. [18] identified OPCAB patients to have significantly less major postoperative complications. Puskas and colleagues [9] compared 14,766 patients and found that OPCAB is associated with a lower operative mortality and disproportionately benefits high-risk patients.

Very similar to our report, Sharoni et al. recently evaluated 353 patients with an EF $<35\%$ who either underwent OPCAB ($n = 144$) or on-pump ($n = 209$) coronary artery bypass surgery. Both groups did not differ significantly in regard to major postoperative complication rates or mortality suggesting OPCAB to be applicable for patients with impaired left ventricular function. The authors showed an increase in the use of OPCAB over time, without any impact on morbidity or mortality [19]. This is in line with our results and is well comparable to the evolution of OPCAB as the standard of care at many institutions.

Cardiopulmonary bypass may have a damaging effect on the myocardium. Pathophysiologically, this is reflected by an extended degree of inflammatory response [4] and myocardial injury [20, 21], especially if the CPB run is prolonged [10]. The activation of numerous inflammatory mediators may compromise myocardial performance, particularly if the LV function is already severely impaired [22, 23]. Moreover, due to the transient change of the ventricular geometry and the ischemic cardiac arrest during CPB, the coronary collateral-flow supplying ischemic areas of the myocardium, is limited [23]. Therefore, OPCAB may be beneficial for patients with severely decreased EF

with regards to an overall better cardiac recovery and straighter postoperative course.

A standardized OPCAB approach in high-risk patients does not come at a price of less complete revascularization. This is an important aspect that has been highlighted to be a crucial predictor for the long-term outcome [7, 8], but has also been reported to be a major argument against OPCAB [11, 24, 25]. Our findings are supported by Puskas et al. [8] who demonstrated feasibility of complete revascularization using OPCAB.

Due to its retrospective nature and non-randomized design, all established disadvantages apply. Although balancing scores constitute the most rigorous methods available for apples-to-apples investigation of causal effects on outcome in the retrospective, nonrandomized setting, they are not equal to randomized clinical trials and they cannot account for unknown variables affecting outcome that are not correlated strongly with measured variables [26]. In addition, our results lack the force of numbers and certainly a higher level of significance may have been achieved, had we had a larger patient cohort to analyze. Although the total number of diseased vessels was included in the propensity adjustment, a certain bias may apply, since OPCAB patients had significantly less diseased vessels. Finally, the study period was quite long with most CABG patients being from the early part of the study, whereas the major part of OPCAB patients was from the later part of the study period.

In conclusion, OPCAB in high-risk patients with severely decreased EF is safe and should not deter surgeons from performing the OPCAB approach in this subset of patients. It comes with similar mortality and MACCE, may even benefit the patients in regard to non-cardiac complications and is not at cost of less complete revascularization.

References

1. Christakis GT, Weisel RD, Fremes SE, Ivanov J, David TE, Goldman BS, Salerno TA (1992) Coronary artery bypass grafting in patients with poor ventricular function. Cardiovascular Surgeons of the University of Toronto. *J Thorac Cardiovasc Surg* 103:1083–1091; discussion 1091–1092
2. O'Connor GT, Plume SK, Olmstead EM, Coffin LH, Morton JR, Maloney CT, Nowicki ER, Levy DG, Tryzelaar JF, Hernandez F (1992) Multivariate prediction of in-hospital mortality associated with coronary artery bypass graft surgery. Northern New England Cardiovascular Disease Study Group. *Circulation* 85:2110–2118
3. Patel NC, Deodhar AP, Grayson AD, Pullan DM, Keenan DJ, Hasan R, Fabri BM (2002) Neurological outcomes in coronary surgery: independent effect of avoiding cardiopulmonary bypass. *Ann Thorac Surg* 74:400–405; discussion 405–406
4. Wan IY, Arifi AA, Wan S, Yip JH, Sihoe AD, Thung KH, Wong EM, Yim AP (2004) Beating heart revascularization with or without cardiopulmonary bypass: evaluation of inflammatory response in a prospective randomized study. *J Thorac Cardiovasc Surg* 127:1624–1631
5. Neshar N, Frolkis I, Vardi M, Sheinberg N, Bakir I, Caselman F, Pevni D, Ben-Gal Y, Sharony R, Bolotin G, Loberman D, Uretzky G, Weinbroum AA (2006) Higher levels of serum cytokines and myocardial tissue markers during on-pump versus off-pump coronary artery bypass surgery. *J Card Surg* 21:395–402
6. Hannan EL, Wu C, Smith CR, Higgins RS, Carlson RE, Culliford AT, Gold JP, Jones RH (2007) Off-pump versus on-pump coronary artery bypass graft surgery: differences in short-term outcomes and in long-term mortality and need for subsequent revascularization. *Circulation* 116:1145–1152
7. Lattouf OM, Thourani VH, Kilgo PD, Halkos ME, Baio KT, Myung R, Cooper WA, Guyton RA, Puskas JD (2008) Influence of on-pump versus off-pump techniques and completeness of revascularization on long-term survival after coronary artery bypass. *Ann Thorac Surg* 86:797–805
8. Puskas JD, Williams WH, Duke PG, Staples JR, Glas KE, Marshall JJ, Leimbach M, Huber P, Garas S, Sammons BH, McCall SA, Petersen RJ, Bailey DE, Chu H, Mahoney EM, Weintraub WS, Guyton RA (2003) Off-pump coronary artery bypass grafting provides complete revascularization with reduced myocardial injury, transfusion requirements, and length of stay: a prospective randomized comparison of two hundred unselected patients undergoing off-pump versus conventional coronary artery bypass grafting. *J Thorac Cardiovasc Surg* 125:797–808
9. Puskas JD, Thourani VH, Kilgo P, Cooper W, Vassiliades T, Vega JD, Morris C, Chen E, Schmotzer BJ, Guyton RA, Lattouf OM (2009) Off-pump coronary artery bypass disproportionately benefits high-risk patients. *Ann Thorac Surg* 88:1142–1147
10. Moshkovitz Y, Sternik L, Paz Y, Gurevitch J, Feinberg MS, Smolinsky AK, Mohr R (1997) Primary coronary artery bypass grafting without cardiopulmonary bypass in impaired left ventricular function. *Ann Thorac Surg* 63:S44–S47
11. Chamberlain MH, Ascione R, Reeves BC, Angelini GD (2002) Evaluation of the effectiveness of off-pump coronary artery bypass grafting in high-risk patients: an observational study. *Ann Thorac Surg* 73:1866–1873
12. Shennib H, Endo M, Benhamed O, Morin JF (2002) Surgical revascularization in patients with poor left ventricular function: on- or off-pump? *Ann Thorac Surg* 74:S1344–S1347
13. Arom KV, Flavin TF, Emery RW, Kshetry VR, Petersen RJ, Janey PA (2000) Is low ejection fraction safe for off-pump coronary bypass operation? *Ann Thorac Surg* 70:1021–1025
14. Emmert MY, Salzberg SP, Seifert B, Schurr UP, Hoerstrup SP, Reuthebuch O, Genoni M (2010) Routine off-pump coronary artery bypass grafting is safe and feasible in high-risk patients with left main disease. *Ann Thorac Surg* 89:1125–1130
15. Kumazaki S, Koyama J, Aizawa K, Kasai H, Koshikawa M, Izawa A, Tomita T, Takahashi M, Ikeda U (2010) Effect of graft adaptation of the internal mammary artery on longitudinal phasic blood flow velocity characteristics after surgery. *Heart Vessels* 25:515–521
16. Shroyer AL, Grover FL, Hattler B, Collins JF, McDonald GO, Kozora E, Lucke JC, Baltz JH, Novitzky D (2009) On-pump versus off-pump coronary-artery bypass surgery. *N Engl J Med* 361:1827–1837
17. Stamou SC, Jablonski KA, Hill PC, Bafi AS, Boyce SW, Corso PJ (2005) Coronary revascularization without cardiopulmonary bypass versus the conventional approach in high-risk patients. *Ann Thorac Surg* 79:552–557
18. Al-Ruzzeh S, Nakamura K, Athanasiou T, Modine T, George S, Yacoub M, Ilsley C, Amrani M (2003) Does off-pump coronary artery bypass (OPCAB) surgery improve the outcome in high-risk patients?: a comparative study of 1398 high-risk patients. *Eur J Cardiothorac Surg* 23:50–55

19. Sharoni E, Song HK, Peterson RJ, Guyton RA, Puskas JD (2006) Off pump coronary artery bypass surgery for significant left ventricular dysfunction: safety, feasibility, and trends in methodology over time—an early experience. *Heart* 92:499–502
20. Koh TW, Carr-White GS, DeSouza AC, Ferdinand FD, Hooper J, Kemp M, Gibson DG, Pepper JR (1999) Intraoperative cardiac troponin T release and lactate metabolism during coronary artery surgery: comparison of beating heart with conventional coronary artery surgery with cardiopulmonary bypass. *Heart* 81:495–500
21. Krejca M, Skiba J, Szmagała P, Gburek T, Bochenek A (1999) Cardiac troponin T release during coronary surgery using intermittent cross-clamp with fibrillation, on-pump and off-pump beating heart. *Eur J Cardiothorac Surg* 16:337–341
22. Rastan AJ, Eckenstein JJ, Hentschel B, Funkat AK, Gummert JF, Doll N, Walther T, Falk V, Mohr FW (2006) Emergency coronary artery bypass graft surgery for acute coronary syndrome: beating heart versus conventional cardioplegic cardiac arrest strategies. *Circulation* 114:1477–1485
23. Stamou SC, Corso PJ (2001) Coronary revascularization without cardiopulmonary bypass in high-risk patients: a route to the future. *Ann Thorac Surg* 71:1056–1061
24. Thomas GN, Martinez EC, Woitek F, Emmert MY, Sakaguchi H, Muecke S, Lee CN, Kofidis T (2009) Off-pump coronary bypass grafting is safe and efficient in patients with left main disease and higher EuroScore. *Eur J Cardiothorac Surg* 36:616–620
25. Yeatman M, Caputo M, Ascione R, Ciulli F, Angelini GD (2001) Off-pump coronary artery bypass surgery for critical left main stem disease: safety, efficacy and outcome. *Eur J Cardiothorac Surg* 19:239–244
26. Blackstone EH (2002) Comparing apples and oranges. *J Thorac Cardiovasc Surg* 123:8–15